

Calculating energy shifts in terms of phase shifts

Zhenhua Yu,^a Gordon Baym,^{a,b} and C. J. Pethick,^{a,c}

^a*The Niels Bohr International Academy, The Niels Bohr Institute,
Blegdamsvej 17, DK-2100 Copenhagen Ø, Denmark*

^b*Department of Physics, University of Illinois, 1110 W. Green Street, Urbana, IL 61801*

^c*NORDITA, Roslagstullsbacken 23, SE-10691 Stockholm, Sweden*

(Dated: May 28, 2010)

To clarify the relation of energy shifts to scattering phase shifts in one-body and many-body problems, we examine their relation in a number of different situations. We derive, for a particle in a container of arbitrary shape with a short-range scattering center, a general result for the energy eigenvalues in terms of the s-wave scattering phase shift and the eigenstates in the absence of the scatterer. We show that, while the energy shifts for a spherical container are proportional to the phase shift over large ranges, those for a cubic container have a more complicated behavior. We connect our result to the description of energy shifts in terms of the scattering T-matrix. The general relation is extended to problems of particles in traps with smoothly varying potentials, including, e.g., the interaction of a small neutral atom with a Rydberg atom. We then consider the many-body problem for particles with a two-body interaction and show that the energy change due to the interaction is proportional to an average of a generalized phase shift that includes the effects of the medium. Finally, we discuss why, even though individual energy levels are very sensitive to boundary conditions, the energy of a many-body system is not.

PACS numbers: 03.65.Nk, 67.85.-d, 31.15.-p

I. INTRODUCTION

The art of calculating energies of many-particle systems in terms of scattering phase shifts has a long history, dating back to Lenz [1], who derived the low-energy effective interaction between particles in terms of the scattering length for two-body scattering, and to Beth and Uhlenbeck [2], who expressed the second virial coefficient for a classical gas in terms of the phase shifts for two-body scattering. Such a procedure is convenient, since it separates the physics of the short-range scattering process from that of single-particle motion in the external potential confining the particles. There are many variants of the formalism, including the pseudopotential approach of Huang and Yang [3].

Further applications, particularly to systems in finite geometry, include the early example of calculation of energy shifts of electronic states due to impurities in a metal [4], and studies of ultracold atomic gases confined in traps. Yet further examples are calculations of Rydberg molecules [5], the calculation of the energy levels of two interacting particles in a harmonic trap [6, 7], and the construction of generalizations of the Gross-Pitaevskii equation to include interactions beyond the usual pseudopotential proportional to the s-wave scattering length between the particles [8, 9]. From a quite different perspective, Lüscher [10] derived a formula for the energies of bound states of two particles in a box (with periodic boundary conditions) that is enjoying use in particle physics to extract scattering properties of elementary particles from lattice calculations of energies [11].

Our primary aim in this paper is to clarify the relation between energy shifts arising from interparticle interactions and the scattering phase shifts. We begin by con-

sidering the energy levels of a particle in a spherical box in the presence of a scatterer at the center of the box. For simplicity, we focus on energies low enough that only s-wave scattering from the scatterer need be taken into account. In the presence of a short-range scatterer, the energy shifts are linear in the s-wave phase shift, δ , and not a trigonometric function of δ , such as the real part of the scattering amplitude. However, in the case of a particle in a more generally shaped container, with more complicated boundary conditions on the wave functions, we find that the energy shifts depend implicitly on $\tan \delta$ (see Eq. (14)). In particular, as we shall see explicitly for a particle in a cubic container, while the energy shifts are approximately linear in δ for small δ , the slope of the shift varies from state to state. In the many body problem, as we illustrate with a simplified separable interaction, a delta-shell potential, the energy shift is linear in an effective scattering phase shift that takes into account effects of the medium.

We begin, in Sec. II, by reviewing the energy shifts of a particle in a spherical container. We then go on in Sec. III to derive a general expression for the energy of a single particle moving in a container of arbitrary shape in the presence of a scattering center. The energy is obtained as an implicit function of the s-wave phase shift of the scatterer and the eigenfunctions of the one-particle problem in the absence of the scatterer. We apply the general result to the case of a cubic container in Sec. IV. In Sec. V we derive results for the cubic container from a T-matrix approach, and in Sec. VI describe applications to situations such as atoms in traps and Rydberg atoms in which the particle moves in a potential that varies smoothly in space, rather than in a container with infinitely repulsive walls. Section VII considers applica-

tions to the many-body problem, and discusses how the formalism may be generalized to take into account the effects of the medium on two-body scattering. We take $\hbar = 1$ throughout.

II. ENERGY SHIFTS IN A SPHERE

To set the stage, we first review the relation of the energy shifts and s-wave scattering phase shifts for the familiar problem of a particle of mass M interacting with a short range potential $v(\mathbf{r})$ in a hard walled spherical container whose radius, R , is much larger than the range, b , of the potential. At low energies, scattering is predominantly s-wave, even if the potential is not central, and may be parametrized in terms of the s-wave phase shift, δ . Consequently, at low energies, eigenstates of the energy are also eigenstates of the angular momentum, and we shall characterize states by the value of the angular momentum. In this limit, states other than s-wave ones are unaffected by the scatterer. For $v(\mathbf{r}) = 0$, the single-particle eigenenergies for s-wave states are $E_\nu = k_\nu^2/2M$, with $k_\nu = \nu\pi/R$, and $\nu = 1, 2, 3, \dots$, with eigenfunctions

$$\psi_\nu(\mathbf{r}) = \frac{1}{\sqrt{2\pi R}} \frac{\sin(k_\nu r)}{r}. \quad (1)$$

In the presence of the scatterer, the eigenfunctions at points beyond the range of the potential become

$$\psi'_\nu(\mathbf{r}) = A_1 \frac{\sin(k'_\nu r + \delta(k))}{r}, \quad (2)$$

where A_1 is a normalization constant. The hard wall boundary condition implies that

$$k'_\nu R + \delta(k'_\nu) = \nu\pi, \quad (3)$$

or

$$\cot(k'_\nu R) = -\cot \delta(k'_\nu). \quad (4)$$

The shift in the wavevector, $\Delta k_\nu \equiv k'_\nu - k_\nu = -\delta(k'_\nu)/R$, is linear in the phase shift. Without loss of generality we may take δ to lie in the range $-\pi/2$ to $\pi/2$, since an increase of δ by π takes one from a given state in the sphere to the next higher state. Since beyond the range of the potential $\psi'_\nu(\mathbf{r})$ is a solution of the Schrödinger equation with zero potential, the corresponding eigenenergies are $E_\nu = k_\nu^2/2M$, and the shifts in the energies due to the interaction are

$$\Delta E_\nu = \frac{k_\nu'^2 - k_\nu^2}{2M} = -\frac{2\pi\delta(k'_\nu)}{Mk_\nu} \left(1 - \frac{\delta(k'_\nu)}{2k_\nu R}\right) |\psi_\nu(0)|^2. \quad (5)$$

The term quadratic in δ may be neglected if $\delta \ll 1$, since $k_\nu R \gtrsim 1$. Even for resonant scattering, $\delta \sim \pi/2$, the quadratic term may be neglected for states with many nodes ($k_\nu R \gg 1$). The energy shift is then given by

$$\Delta E_\nu \approx -\frac{2\pi\delta(k'_\nu)}{Mk_\nu} |\psi(0)|^2. \quad (6)$$

Thus in these general situations, the energy shift is linear in the phase shift (and not, e.g., the real part of the scattering amplitude $\sim e^{i\delta} \sin \delta$). The energy of a state must be determined self-consistently, since δ depends on energy, but if $R^{-1}(|d\delta/dk|) \ll 1$ one may replace k'_ν by the wave number of the state in the absence of the scatterer, k_ν .

In the limit $k \rightarrow 0$, away from a scattering resonance, $\delta \rightarrow -ka_s$, with a_s the s-wave scattering length. Then Eq. (6) gives

$$\Delta E(k) = \frac{2\pi a_s}{M} |\psi(0)|^2. \quad (7)$$

III. S-WAVE SCATTERING IN AN ARBITRARY CONTAINER

We turn now to establishing, for a particle of mass M interacting with a short range central potential in a container of arbitrary shape, the relation between the energy shifts of the particle and the s-wave scattering phase shift. We consider a particle interacting with a scatterer located at some point within the container, which we take to be $\mathbf{r} = 0$. We shall take the potential to be spherically symmetric and short-range, vanishing for $r > b$, and we assume that the characteristic dimension of the container, λ , is much larger than the range of the potential, b . We shall consider energies, E , sufficiently low that only s-wave scattering is important. We denote the eigenfunction $\psi_k(\mathbf{r})$ for energy $E = k^2/2M$ by $\psi_k^<(\mathbf{r})$ for $r < b$ and $\psi_k^>(\mathbf{r})$ for $r > b$. For distances $r \ll \lambda$, $\psi_k^>(\mathbf{r})$ may be expanded in terms of spherical waves as

$$\psi_k^>(\mathbf{r}) = A_2 \left[\frac{\sin(kr)}{kr} + \tan \delta \frac{\cos(kr)}{kr} + \sum_{l=1}^{\infty} \sum_{m=-l}^l \alpha_{lm} j_l(kr) Y_{lm}(\theta, \phi) \right], \quad (8)$$

where j_l is the spherical Bessel function, the α_{lm} are expansion coefficients, and A_2 is a normalization constant. In the sum, terms proportional to the spherical Neumann function n_l , which is singular for $r \rightarrow 0$, have been omitted, since we assume that only s-wave scattering is important, and consequently the wave function for other partial waves must be finite for $r \rightarrow 0$. Thus asymptotically, in the region $b < r \ll 1/k$,

$$\psi_k^>(\mathbf{r}) = A_2 \left[\frac{\tan \delta}{kr} + 1 + \mathcal{O}(kr) \right]. \quad (9)$$

To derive an expression for the energy of a state, we now find a second expression for the wave function, by expressing it in terms of the single particle eigenstates $\{\phi_n(\mathbf{r})\}$ for $v(\mathbf{r}) = 0$, with energy E_n as follows. The single particle Green function for $v(\mathbf{r}) = 0$ describing

propagation of a particle from the origin to $\mathbf{r}$ is given by

$$G_k(\mathbf{r}, 0) = \sum_n \frac{\phi_n(\mathbf{r})\phi_n^*(0)}{E_n - E}. \quad (10)$$

This satisfies the equation,

$$\frac{1}{2M}(\nabla^2 + k^2)G_k(\mathbf{r}, 0) = -\delta(\mathbf{r}), \quad (11)$$

which shows that $G_k(\mathbf{r}, 0)$ satisfies the Schrödinger equation for $\psi_k^>(\mathbf{r})$ when $\mathbf{r} \neq 0$. Thus outside the range of the potential, $G_k(\mathbf{r}, 0)$ satisfies the Schrödinger equation and in addition satisfies the same boundary conditions as the wave functions ϕ_n at the walls of the container for any boundary condition that is linear in the wave function. Equation (11) implies that $G_k(\mathbf{r}, 0) = M/2\pi r$ for $kr \ll 1$. One can in fact show that for $r > b$

$$\psi_k^>(\mathbf{r}) = \frac{2\pi A_2 \tan \delta}{Mk} G_k(\mathbf{r}, 0), \quad (12)$$

by following Ref. [10]. The function

$$\chi_k(\mathbf{r}) = \psi_k^>(\mathbf{r}) - \frac{2\pi A_2 \tan \delta}{Mk} G_k(\mathbf{r}, 0) \quad (13)$$

is regular everywhere inside the container and is an eigenfunction with eigenenergy E for $v(\mathbf{r}) = 0$. However, unless $E = E_n$, this is impossible, since the ϕ_n form a complete set, and therefore $\chi_k(\mathbf{r}) = 0$. (For discussion of the more complicated case when $E = E_n$, see Ref. [10].)

Matching the singular and nonsingular parts in Eqs. (9) and (12) for $kr \ll 1$, we obtain an expression for the energy of the state:

$$\frac{k}{4\pi} \cot \delta(k) = \lim_{r \rightarrow 0} \left[\sum_n \frac{1}{2M} \frac{\phi_n(\mathbf{r})\phi_n^*(0)}{E_n - E} - \frac{1}{4\pi r} \right], \quad (14)$$

since non s-wave contributions vanish in the limit $r \rightarrow 0$.

Equation (14) is one of the main results of the paper, and determines the eigenenergies in the presence of the potential. When the potential strength tends to zero, E approaches an unperturbed energy $\mathcal{E}$; accordingly Eqs. (10) and (12) implies that $\psi_k(\mathbf{r})$ approaches the state

$$\phi_{\mathcal{E}}^s(\mathbf{r}) = C \sum_{E_n = \mathcal{E}} \phi_n(\mathbf{r})\phi_n^*(0), \quad (15)$$

where the sum is over all possible degenerate states of energy $\mathcal{E}$, and $C = (\sum_{E_n = \mathcal{E}} |\phi_n(0)|^2)^{-1/2}$ is the normalization constant. By construction, $\phi_{\mathcal{E}}^s(\mathbf{r})$ has a nonzero s-wave component, i.e., $\phi_{\mathcal{E}}^s(0) \neq 0$ if not all $\phi_n^*(0)$ equal zero. The remaining states $\phi_{\mathcal{E}}^{\bar{s}}(\mathbf{r})$ in the subspace spanned by the states with $E_n = \mathcal{E}$ vanish at $\mathbf{r} = 0$, as we now show. Since these states are energy eigenstates in the absence of the scatterer, we may write $\phi_{\mathcal{E}}^{\bar{s}}(\mathbf{r}) = \sum_{E_n = \mathcal{E}} c_n \phi_n(\mathbf{r})$. We now evaluate the integral $\int d\mathbf{r} (\phi_{\mathcal{E}}^s(\mathbf{r}))^* \phi_{\mathcal{E}}^{\bar{s}}(\mathbf{r})$, which by orthogonality is zero

if $s \neq \bar{s}$. Inserting Eq. (15) and the expansion for $\phi_{\mathcal{E}}^s(\mathbf{r})$ in the integral, one finds $\int d\mathbf{r} (\phi_{\mathcal{E}}^s(\mathbf{r}))^* \phi_{\mathcal{E}}^{\bar{s}}(\mathbf{r}) = C \sum_{E_n = \mathcal{E}} c_n \phi_n(0) = C \phi_{\mathcal{E}}^s(0) = 0$. Therefore the $\phi_{\mathcal{E}}^{\bar{s}}(\mathbf{r})$ for $s \neq \bar{s}$ vanish at the origin and these states will be unaffected when the scatterer at $\mathbf{r} = 0$ is introduced.

For $k \rightarrow 0$, away from a scattering resonance, the left side of Eq. (14) becomes $-1/4\pi a_s$, and so for a_s small compared with a characteristic dimension of the container, E must be very close to an unperturbed eigenenergy E_n . Changing the order of taking the limit $r \rightarrow 0$ and summation in Eq. (14), we obtain

$$\Delta E = E - E_n = \frac{2\pi a_s}{M} |\phi_{E_0}^s(0)|^2; \quad (16)$$

thus the result (7) found in a sphere holds for more general geometry.

We now summarize the salient results of this section. When the energies of states of a particle in the container in the absence of the scatterer are degenerate, one may construct one state, Eq. (15), that does not vanish at $r = 0$, while all other orthogonal states in the degenerate manifold vanish at $r = 0$. For weak scattering, $|\delta(k)| \ll \pi$, the energy shift of the first state is given by Eq. (16) while the shift of the other states vanishes. In our derivation of these results, we have taken into account degeneracy of the energy levels in the absence of the scatterer. For stronger scattering, it is necessary to solve Eq. (14).

The energy shift ΔE is reproduced by the expectation value of the effective interaction [12],

$$V_{\text{eff}}(\mathbf{r}) = \frac{2\pi a_s}{M} \delta(\mathbf{r}), \quad (17)$$

in the *unperturbed* state, ϕ_n . In terms of the interaction of two particles of mass m , the coefficient of the effective interaction is $4\pi a_s/m$.

One should note that the effective interaction (17) is not useful for calculating the perturbed wave function, $\psi_>(r)$, outside the range of the potential in terms of a boundary condition at the origin. Here one needs rather to employ the pseudopotential [3]

$$V_{\text{ps}}(\mathbf{r}) = -\frac{2\pi \tan \delta}{Mk} \delta(\mathbf{r}) \frac{\partial}{\partial r} r \quad (18)$$

in the Schrödinger equation

$$\left(-\frac{\nabla^2}{2M} + V_{\text{ps}}(\mathbf{r}) \right) \psi_>(r) = \frac{k^2}{2M} \psi_>(r). \quad (19)$$

The solution is clearly Eq. (8). The present discussion is readily generalized to higher angular momentum, with a pseudopotential involving higher derivatives of the wave function [13].

IV. ENERGY SHIFTS IN A CUBIC BOX

We now apply Eq. (14) to the specific example of scattering by a spherically symmetric potential centered in a

cubic box of side L , with periodic boundary conditions. This geometry, used in the calculations of Refs. [10, 11], is more difficult to calculate analytically than in a spherical container, since the boundary is incompatible with the rotational symmetry of the central potential. As one sees explicitly by direct construction of the solutions of the Schrödinger equation, all partial waves are mixed in determining the eigenenergies [10].

In the absence of the scatterer, the states are simply

$$\phi_{\mathbf{p}}(\mathbf{r}) = \frac{1}{\sqrt{L^3}} e^{i\mathbf{p}\cdot\mathbf{r}}, \quad (20)$$

where $\mathbf{p} = \{p_x, p_y, p_z\} = (2\pi/L)\{l_x, l_y, l_z\}$ with $l_x, l_y, l_z = 0, \pm 1, \pm 2, \dots$. As before, we consider only the s-wave contribution to the interaction, and from Eq. (14) find [10]

$$\frac{k}{4\pi} \cot \delta(k) = \lim_{r \rightarrow 0} \left[\frac{1}{L^3} \sum_{\mathbf{p}} \frac{e^{i\mathbf{p}\cdot\mathbf{r}}}{p^2 - k^2} - \frac{1}{4\pi r} \right]. \quad (21)$$

A relation similar to Eq. (21) is more useful for simulations on a lattice [11]:

$$\frac{k}{4\pi} \cot \delta(k) = \frac{1}{L^3} \sum_{|\mathbf{p}| < \Lambda} \frac{1}{p^2 - k^2} - \frac{\Lambda}{4\pi}, \quad (22)$$

where Λ is the momentum cutoff on the lattice. To derive Eq. (22), we note that in the limit $r \rightarrow 0$ the terms in the sum in Eq. (21) for $|\mathbf{p}| > \Lambda$ and $k \ll \Lambda$ can be written as

$$\begin{aligned} \frac{1}{L^3} \sum_{|\mathbf{p}| > \Lambda} \frac{e^{i\mathbf{p}\cdot\mathbf{r}}}{p^2} &= \int_0^\infty \frac{d^3p}{(2\pi)^3} \frac{e^{i\mathbf{p}\cdot\mathbf{r}}}{p^2} - \int_0^\Lambda \frac{d^3p}{(2\pi)^3} \frac{e^{i\mathbf{p}\cdot\mathbf{r}}}{p^2} \\ &= \frac{1}{4\pi r} - \frac{\Lambda}{4\pi}. \end{aligned} \quad (23)$$

For small δ , the energy shift obeys Eq. (16). Whereas the shift in k for a sphere is linear in $-\delta$, for a cube it is only approximately linear in $-\delta$, as we see from Fig. (1), which gives a plot of k (panel (a)) and E (panel (b)) vs. $-\delta$ for the first few eigenstates, calculated from Eq. (22). When δ is small (except in the ground state),

$$k'_l L/2\pi \approx k_l L/2\pi - \alpha_{k_l} \delta, \quad (24)$$

where $E = k_l^2/2M$ is the eigenenergy in the absence of the scatterer, and from Eq. (22), $\alpha_{k_l} = D_{k_l}/(k_l L)^2$, with $D_{k_l} = \sum_{\mathbf{p}} \delta_{p^2, k_l^2}$ the degeneracy of the level E . Therefore $\Delta k'_l L/2\pi \approx -\delta D_{k_l}/(k_l L)^2$ for small δ , where $\Delta k'_l = k'_l - k_l$. The linear relation is valid only for small δ , but with a slope which varies with the eigenstate of the cubic box.

The different dependences of k and E on δ for cubic and spherical boxes are due solely to the different boundary conditions. The information on the boundary conditions is encoded in Eq. (14) through the unperturbed eigenvalues and eigenfunctions, while the effect of the short-range potential is taken into account via δ ; in an arbitrary container, the energy change is a function of $\cot \delta$ but the exact form of the function is determined by the boundary conditions.




FIG. 1: (a) Wavenumbers k and (b) eigenenergies E vs. $-\delta$ in a cubic box. The curves are calculated from Eq. (22) with $\Lambda = 200\pi/L$.

V. T-MATRIX APPROACH

With a view to generalizing results to situations other than dilute gases, where binary collisions are the most important interaction process, it is useful to derive the result (22) in terms of the T-matrix. The formalism may then be applied to calculate properties of interacting many-body systems, when the elementary excitations are quite different from free particles, e.g. liquid ^3He at low temperatures. Equation (14) was derived by inspecting the Schrödinger equation with boundary conditions at both small and large r . The T-matrix formalism we give below is equivalent to solving the Schrödinger equation in the one-body problem and its analogue in the many-body problem to be discussed later provides a convenient way to treat the interparticle interactions.

The T-matrix is defined symbolically as

$$T(E) = v + v \frac{1}{E - H_0} T(E) = v + v \frac{1}{E - H} v, \quad (25)$$

where H_0 is the free particle Hamiltonian, v is the potential operator, $H = H_0 + v$, $(E - H)^{-1}$ is the particle propagator, and E the energy; this equation shows that the T-matrix has poles in E at the eigenenergies of the full Hamiltonian. To derive Eq. (22) we rewrite (25) as:

$$T^{-1}(E) = v^{-1} - \frac{1}{E - H_0}; \quad (26)$$

with the on-shell T-matrix, at $E = k^2/2M + i\eta$, given by

$$T^{-1}(k^2/2M + i\eta) = -\frac{Mk}{2\pi} (\cot \delta(k) - i). \quad (27)$$

For simplicity we assume a short ranged contact interaction, $U\delta(\mathbf{r})$ with a momentum cutoff, Λ . Then in free space Eq. (26) becomes,

$$T^{-1}(E) = U^{-1} - \int_0^\Lambda \frac{d^3p}{(2\pi)^3} \frac{1}{E - p^2/2M}. \quad (28)$$

Similarly in the cubic box,

$$T_{\text{box}}^{-1}(E) = U^{-1} - \frac{1}{L^3} \sum_{|\mathbf{p}| < \Lambda} \frac{1}{E - p^2/2M}. \quad (29)$$

Subtracting Eq. (29) from Eq. (28), evaluating both sides at $E = k^2/2M + i\eta$, where $k^2/2M$ is now an eigenenergy in the box, using Eq. (27) and the vanishing of T_{box}^{-1} at $k^2/2M$, we find

$$\begin{aligned} \frac{k}{4\pi} (\cot \delta(k) - i) &= \int_0^\Lambda \frac{d^3p}{(2\pi)^3} \frac{1}{k^2 + i\eta - p^2} \\ &\quad - \frac{1}{L^3} \sum_{|\mathbf{p}| < \Lambda} \frac{1}{k^2 + i\eta - p^2}. \end{aligned} \quad (30)$$

The integral on the right in the limit $k \ll \Lambda$ is $-(\Lambda + ik)/4\pi$. Dropping the $i\eta$ in the sum, we recover Eq. (22).

To see that the general formalism leads to $\Delta k_n R = -\delta$ for a particle in a spherical box (cf. Eq. (3)), we apply the T-matrix formalism. For s-waves in a spherical box, the analog of Eq. (29) is

$$T_{\text{sph}}^{-1}(E) = U^{-1} - \frac{1}{2\pi R} \sum_{p < \pi\nu_{\text{max}}/R} \frac{p^2}{E - p^2/2M}, \quad (31)$$

where the p 's take on the values $\nu\pi/R$, with $\nu = 1, 2, 3, \dots, \nu_{\text{max}}$. By the same argument as for the cubic box, we derive the result

$$k \cot \delta = -\frac{2}{\pi} \Lambda + 2\nu_{\text{max}} - \frac{2k^2}{R} \sum_{p < \pi\nu_{\text{max}}/R} \frac{1}{k^2 - p^2}. \quad (32)$$

Using

$$\sum_n \frac{1}{z - n\pi} = \cot z, \quad (33)$$

we obtain

$$\cot \delta(k) = -\cot(kR) + \frac{1}{kR} - \frac{2}{kR} \left(\frac{\Lambda R}{\pi} - \nu_{\text{max}} \right). \quad (34)$$

Thus to recover the correct relation, $\Delta k R = -\delta$, we need to choose the continuum cutoff Λ to be $\pi(\nu_{\text{max}} + 1/2)/R$, i.e., half way between successive ν values. This subtlety shares a similar origin as in the transformation of a sum of $1/n$ into an integral leading to Euler's constant $\gamma \approx 0.577$. An alternative way of dealing with this problem is to introduce a smooth, rather than a sharp, cutoff.

To further illustrate the T-matrix approach, we consider the delta shell potential

$$v(r) = (v_0/4\pi r_0^2) \delta(r - r_0), \quad (35)$$

which has the advantage that T-matrix can be constructed explicitly without the need for a cutoff. In the continuum limit, where we use as a basis the s-wave functions,

$$\langle \mathbf{r} | k \rangle = \frac{\sin(kr)}{kr}, \quad (36)$$

the T-matrix equation (25) becomes,

$$T_{kk'}(E) = v_{kk'} + \int \frac{d^3p}{(2\pi)^3} \frac{v_{kp} T_{pk'}(E)}{E - p^2/2M}. \quad (37)$$

The matrix elements of the delta shell potential are separable:

$$v_{kk'} = v_0 \frac{\sin(kr_0)}{kr_0} \frac{\sin(k'r_0)}{k'r_0} \equiv v_0 u_k u_{k'}; \quad (38)$$

thus

$$T_{kk'}^{-1}(E) = v_{kk'}^{-1} \left[1 - v_0 \int_0^\infty \frac{d^3p}{(2\pi)^3} \frac{u_p^2}{E - p^2/2M} \right]. \quad (39)$$

On shell ($k = k'$ and $E = k^2/2M + i\eta$),

$$T_{kk}(k^2/2M + i\eta) = \frac{v_0 \sin^2(kr_0)}{k^2 r_0^2 + iMk v_0 (1 - e^{2ikr_0})/4\pi}. \quad (40)$$

Comparing with Eq. (27) we find with some trigonometry that the s-wave phase shift for the delta-shell potential is given by

$$\cot \delta(k) = -\frac{2\pi k r_0^2}{M v_0} [1 + \cot^2(kr_0)] - \cot(kr_0). \quad (41)$$

Similarly in a sphere of radius R , with the s-wave functions

$$\langle r | n \rangle = \frac{1}{\sqrt{2\pi R}} \frac{\sin(k_n r)}{r} \quad (42)$$

as a basis, where $k_n = n\pi/R$, the T-matrix equation (25) becomes,

$$T_{mn}(E) = v_{mn} + \sum_{l=1}^{\infty} v_{ml} \frac{1}{E - E_l} T_{ln}(E); \quad (43)$$

the matrix elements of the delta-shell potential are now

$$v_{mn} = \frac{v_0}{2\pi R r_0^2} \sin(k_m r_0) \sin(k_n r_0). \quad (44)$$

The solution of Eq. (43) is [14]

$$T_{mn}^{-1}(E) = v_{mn}^{-1} \left[1 - \frac{v_0}{2\pi R r_0^2} \sum_{l=1}^{\infty} \frac{\sin^2(k_l r_0)}{E - k_l^2/2M} \right]. \quad (45)$$

For $E = k^2/2M$,

$$\sum_{l=1}^{\infty} \frac{\sin^2(k_l r_0)}{k^2 - k_l^2} = \frac{R}{2k} \sin^2(kr_0) [\cot(kR) - \cot(kr_0)], \quad (46)$$

so that

$$T_{mn}^{-1} = v_{mn}^{-1} \left[1 - \frac{M v_0}{2\pi k r_0^2} \sin^2(kr_0) [\cot(kR) - \cot(kr_0)] \right]. \quad (47)$$

Comparing Eq. (47) with Eq. (41), we find that at the pole of T_{mn} , at eigenenergy $E = k^2/2M$, the expected result

$$\cot(kR) = -\cot(k); \quad (48)$$

this model calculation directly reproduces Eq. (4) (with $k = k'_\nu$) within the T-matrix formalism for the delta shell potential, avoiding the subtle issue of the cutoff Λ in Eq. (34).

VI. PARTICLES IN TRAPS AND RYDBERG ATOMS

So far we have considered containers in which the potential is zero outside the range of the scatterer. The results may be extended to particles in a potential, $V(\mathbf{r})$, such as in an atomic trap. If the scatterer is at point $\mathbf{r}_s$, the Green function for the particle in the potential satisfies the equation

$$\left[-\frac{1}{2M} (\nabla^2 + k^2) + V(\mathbf{r}) - V(\mathbf{r}_s) \right] G_k(\mathbf{r}) = \delta(\mathbf{r}_s), \quad (49)$$

where $k^2 = 2M[E - V(\mathbf{r}_s)]$. Provided $V(\mathbf{r}) - V(\mathbf{r}_s)$ is small compared with other energy scales in Eq. (49) at distances from the scatterer of the order of the range of the potential, the arguments of Sec. III still hold, with the origin shifted to $\mathbf{r}_s$. Explicitly, the condition is that $|V(\mathbf{r}) - V(\mathbf{r}_s)| \ll 1/Mb^2$ for $|\mathbf{r} - \mathbf{r}_s| \lesssim b$, in addition to the usual one $kb \lesssim 1$, where b is the range of the scattering

potential (not $V(\mathbf{r})$). Equation (14) thus holds for this more general case, provided the ϕ_n are taken to be the eigenfunctions including the effects of the potential $V(\mathbf{r})$.

The general result (14) can be applied to calculate the effective potential between a Rydberg atom and a small neutral atom. In the absence of the neutral atom, the eigenstates $\phi_n(\mathbf{r})$ for the outermost electron are hydrogenic wavefunctions for the Coulomb potential $V_c(r)$ exerted by the ion, where we take the nucleus to be at the origin. We place the neutral atom at position $\mathbf{r}_s$, of magnitude much larger than a_0 (the Bohr radius). The eigenenergies E of the outmost electron of the Rydberg atom are functions of $\mathbf{r}_s$, and thus act as an effective potential between the neutral and Rydberg atoms. As a consequence of the electron's interaction with the neutral atom, the wave function of the electron in the Rydberg atom acquires a form near $\mathbf{r} = \mathbf{r}_s$ that corresponds to an s-wave phase shift $\delta(k)$. Thus from Eq. (14), the effective potential $E(r_s)$ is determined by

$$\frac{k}{4\pi} \cot \delta(k) = \lim_{\mathbf{r} \rightarrow \mathbf{r}_s} \left[\sum_n \frac{1}{2m_e} \frac{\phi_n(\mathbf{r}) \phi_n^*(\mathbf{r}_s)}{E_n - E(\mathbf{r}_s)} - \frac{1}{4\pi |\mathbf{r} - \mathbf{r}_s|} \right], \quad (50)$$

where $k = \sqrt{2m_e[E - V_c(r_s)]}$ is the scattering wavelength and m_e is the electron mass. The sum over n includes both the discrete and continuum states. Solving this equation is an alternative route to the results of Ref. [5] for the interaction of a neutral atom with a Rydberg atom. When applied to the problem of two particles in a harmonic potential interacting via a short-range potential, Eq. (50) reproduces the result of Ref. [6].

VII. ENERGY SHIFTS IN MANY-PARTICLE SYSTEMS

We give now an example that allows one to identify the quantity in the many-body problem that plays the role of the phase shift in two-body scattering, and that shows how the effects of the medium enter the scattering process. This section may be regarded as an illustration of the basic formalism for describing thermodynamic properties of a many-body system in terms of fully renormalized single- and many-particle propagators given by De Dominicis and Martin [15] and reviewed in Ref. [16]. We employ the delta shell model and calculate its effect in a dilute binary fermionic gas within the ladder approximation [17, 18]. We assume that the system contains equal numbers of fermions in internal states $|1\rangle$ and $|2\rangle$ of equal mass m , with chemical potential μ .

Differentiating the grand potential per unit volume, Ω , with respect to the coupling constant v_0 gives

$$\begin{aligned} \frac{\partial \Omega}{\partial v_0} &= -\frac{1}{4\pi r_0^2} \int d^3\rho \delta(\rho - r_0) \langle \psi_1^\dagger(\boldsymbol{\rho}) \psi_2^\dagger(0) \psi_2(0) \psi_1(\boldsymbol{\rho}) \rangle \\ &= \frac{1}{\beta V} \sum_{\omega, \mathbf{p}} e^{\omega\eta} \frac{B(p, \omega)}{1 - v_0 B(p, \omega)}, \end{aligned} \quad (51)$$

where η is a positive infinitesimal and two-particle propagator is

$$B(p, \omega) = -\frac{1}{\beta V} \sum_{z, \mathbf{q}} \frac{u_q^2}{(z - \epsilon_{\mathbf{p}/2 + \mathbf{q}})(\omega - z - \epsilon_{\mathbf{p}/2 - \mathbf{q}})} \\ = \int \frac{d^3 q}{(2\pi)^3} u_q^2 \frac{1 - 2f(\epsilon_{\mathbf{p}/2 + \mathbf{q}})}{\omega + 2\mu - p^2/4m - q^2/m}, \quad (52)$$

$\epsilon_{\mathbf{q}} = q^2/2m - \mu$, ω is a bosonic Matsubara frequency, and z a fermionic one, f is the Fermi distribution function, $\beta = 1/T$ with T the temperature (we set the Boltzmann constant to unity), and V is the volume. Here we approximate the single particle propagators by those for free particles. In the low density limit, at zero temperature, $1 - v_0 B(0, \omega)$ is proportional to the inverse T-matrix for the separable potential and is defined in Eq. (39).

Integration of Eq. (51) with respect to v_0 gives the change in the grand potential due to the interaction:

$$\Delta\Omega = -\frac{1}{\beta} \sum_{\omega} \int \frac{d^3 p}{(2\pi)^3} e^{\omega\eta} \ln[t(p, \omega)/v_0], \quad (53)$$

where

$$t(p, \omega) = \frac{v_0}{1 - v_0 B(p, \omega)} \quad (54)$$

is related to the T-matrix in the medium by $T_{kk'}(p, \omega) = u_k u_{k'} t(p, \omega)$, with $u_k = \sin kr_0/kr_0$.

We convert the sum in Eq. (53) into an integral, choosing the branch cut of the logarithm to lie along the negative real axis. We assume that $1 + mv_0/4\pi r_0 \geq 0$ so that there is no two-body bound state, and that the temperature exceeds the BCS transition temperature. Then the only singularity of $t(\mathbf{p}, \omega)$ is the branch cut along the real ω axis, so that

$$\Delta\Omega = - \int \frac{d^3 p}{(2\pi)^3} \int_{-\infty}^{\infty} \frac{d\omega}{\pi} f_B(\omega) \delta_m(p, \omega), \quad (55)$$

where f_B is the Bose distribution function. The quantity

$$\delta_m(p, \omega) = \arg(t(p, \omega + i\eta)) \quad (56)$$

is the analog of δ in the two body problem (cf. Eq. (27)). Therefore

$$\Delta\Omega = \int \frac{d^3 p_1}{(2\pi)^3} \frac{d^3 p_2}{(2\pi)^3} f(\epsilon_{\mathbf{p}_1}) f(\epsilon_{\mathbf{p}_2}) v_{\text{eff}}(\mathbf{p}_1, \mathbf{p}_2). \quad (57)$$

By the theorem of small increments, small changes in thermodynamic potentials keeping the natural variables appropriate to the chosen potential are the same [19]. Consequently, Eq. (57) gives the change in the energy density for fixed particle number and entropy per unit volume. Equation (57) has the usual form for the change in the energy density for a two-body interaction, with an effective two-body interaction given by

$$v_{\text{eff}}(\mathbf{p}_1, \mathbf{p}_2) = \frac{\delta_m(|\mathbf{p}_1 + \mathbf{p}_2|, \epsilon_{\mathbf{p}_1} + \epsilon_{\mathbf{p}_2})}{\text{Im}B(|\mathbf{p}_1 + \mathbf{p}_2|, \epsilon_{\mathbf{p}_1} + \epsilon_{\mathbf{p}_2} + i\eta)}. \quad (58)$$

In low density limit, $-\text{Im}B(|\mathbf{p}_1 + \mathbf{p}_2|, \epsilon_{\mathbf{p}_1} + \epsilon_{\mathbf{p}_2} + i\eta)/\pi$ reduces to $m|\mathbf{p}_1 - \mathbf{p}_2|/8\pi^2$, the density of states for a pair of atoms with total momentum $|\mathbf{p}_1 + \mathbf{p}_2|$. For small $|\mathbf{p}_1 - \mathbf{p}_2|$, the phase shift is given in terms of the s-wave scattering length by $\delta \rightarrow -|\mathbf{p}_1 - \mathbf{p}_2|a_s/2$. The factor of two arises because in the problem of scattering from a rigid scattering center considered earlier, we worked in terms of k , the momentum of the scattered particle, while in the two-body problem, the corresponding quantity is $|\mathbf{p}_1 - \mathbf{p}_2|/2$, the magnitude of the momentum of one of the particles relative to the center of mass of the pair. The effective interaction in momentum space then reduces to the well-known result $4\pi a_s/m$ for two particles of equal mass (see Eq. (17) and the remarks following).

The many-body calculation above shows that the total energy change is linear in a generalized phase shift that includes the effects of the medium on the scattering process. By contrast, Eq. (14) shows that in an arbitrary container, the energy changes of one-particle eigenstates in the presence of a short-range scatterer are generally not linear in the s-wave phase shift, and one similarly expects the energy eigenstates of two or more particles with binary interactions in a container of arbitrary shape not to be linear in the phase shift for the two-body interaction. How does one account for this difference? The essential difference between the two situations may be seen by noting that the energy of a (non-interacting) Fermi gas in the presence of a central scatterer is a sum of contributions from a large number of individual single-particle energy levels. When the number of particles N is large compared with unity, the dominant contribution to the total energy is insensitive to the behavior of any particular energy level. This is true even though the single-particle energies are not linear functions of δ , but exhibit the undulating behavior shown in Fig. 1. Contributions beyond the leading term are, however, sensitive to the shape of the container and are responsible for, e.g., shell structure in nuclear binding energies. Similarly, in the many-body problem with two-body scattering described above, the total energy is insensitive to the boundary conditions a single-particle state satisfies on the walls of the container.

VIII. CONCLUDING REMARKS

One of the main results of this paper is a generalization of an expression due to Lüscher [10] that expresses energy shifts of states in terms of scattering properties. In particular, we have derived a general expression, Eq. (14) for the shift of the energy of the state of a single particle in a container of any shape external potential when a scatterer of short range is added. In addition, a related result for a particle in a potential that is slowly varying in space was derived, Eq. (50). The numerical results obtained for a particle in a cubic box show that the energies of low-lying states are a rather complicated function of the phase shift. However, the functional form is con-

strained by the fact that when the phase shift is reduced from δ to $\delta - \pi$, the state of the system is transformed into the next higher state for a phase shift δ .

In Sec. VII, we considered the many-body problem and showed that, for a model system of two species of fermion interacting via a two-body potential, the shift in the energy may be expressed as a *linear* function of an effective interaction proportional to a generalized phase shift that takes into account the effect of the medium on the scattering process. The reason for this simple behavior for the many-body problem is that the free energy is a thermodynamic quantity, and this is sensitive mainly to the

volume of the container, but insensitive to the details of the single particle wave functions, which depend on the shape of the container.

Acknowledgements

We are grateful to David Kaplan for calling our attention to the applications of the Lüscher formula in lattice simulations. This research was supported in part by NSF Grant PHY07-01611.

-
- [1] W. Lenz, Z. Phys. **56**, 778 (1929).
 - [2] G. E. Uhlenbeck and E. Beth, Physica **3**, 729 (1936); E. Beth and G. E. Uhlenbeck, Physica **4**, 915 (1937).
 - [3] K. Huang and C. N. Yang, Phys. Rev. **105**, 767 (1957).
 - [4] F. G. Fumi, Philos. Mag. **46**, 1007 (1955).
 - [5] C. H. Greene, A. S. Dickinson, and H. R. Sadeghpour, Phys. Rev. Lett. **85**, 2458 (2000); E. L. Hamilton, C. H. Greene, and H. R. Sadeghpour, J. Phys. B **35**, L199 (2002).
 - [6] T. Busch, B. Englert, K. Rzażewski, and M. Wilkens, Found. Phys. **28**, 549 (1997).
 - [7] D. Blume and C. H. Greene, Phys. Rev. A **65**, 043613 (2002); E. L. Bolda, E. Tiesinga, and P. S. Julienne, Phys. Rev. A **66**, 013403 (2002); B. Borca, D. Blume, and C. H. Greene, New J. of Physics **5**, 111.1 (2003).
 - [8] H. Fu, Y. Wang, and B. Gao, Phys. Rev. A **67**, 053612 (2003).
 - [9] A. Collin, P. Massignan, and C. J. Pethick, Phys. Rev. A **75**, 013615 (2007).
 - [10] M. Lüscher, Nucl. Phys. B **354**, 531 (1991); also, Com. Math. Phys. **105**, 153 (1986).
 - [11] S. R. Beane, P. F. Bedaque, A. Parreño, and M. J. Savage, Phys. Lett. B **585**, 106 (2004); Nucl. Phys A **747**, 55 (2005).
 - [12] A. J. Leggett, *Quantum Liquids: Bose Condensation and Cooper Pairing in Condensed-Matter Systems*, (Oxford Univ. Press, Oxford, 2007).
 - [13] R. Roth and H. Feldmeier, Phys. Rev. A **64**, 043603 (2001).
 - [14] The delta shell potential is closely related to the contact pseudopotential $U\delta(\mathbf{r})$ with cutoff Λ , with the identification $U \sim v_0$ and $\Lambda \sim 1/r_0$. The factor $(\sin k_l r_0)^2$ in Eq. (45) at small k_r is $\sim k_l^2$, as in Eq. (31); however the k_l^2 behavior flattens out at $k_l \sim 1/r_0$, regularizing the sum.
 - [15] C. T. De Dominicis and P. C. Martin, J. Math. Phys. **5**, 14, 31 (1964).
 - [16] C. Bloch, *Studies in Statistical Mechanics*, Vol. III, ed. J. de Boer and G. E. Uhlenbeck, (North Holland, Amsterdam, 1965) p. 1.
 - [17] P. Nozières and S. Schmitt-Rink, J. Low Temp. Phys. **59**, 195 (1985).
 - [18] Similar arguments have been given for a contact interaction by C. J. Pethick and H. Smith, *Bose-Einstein condensation in dilute gases*, 2nd edition, (Cambridge Univ. Press, Cambridge, 2008), Sec. 17.4. The analogous results for particle-hole scattering were used in C. J. Pethick and G. M. Carneiro, Phys. Rev. A **7**, 304 (1973).
 - [19] L. D. Landau and E. M. Lifshitz, *Statistical Physics*, Vol. I, Third edition, (New York, Pergamon, 1980), §15.